北京景山学校 Name: Grade /100

Maths - Pre-Calc. - Senior 1+ - REVIEW TEST 1 Nov.15 (60 min.) - p.1/3


I – **Linear Programming** : [30 pts] A publishing company is producing two kinds of software A and B on DVDs. It uses two machines for the production : one for the disk burning and one for the packaging.

Let x be the number of DVDs of A type and y be the number of DVDs of B type.

The burning machine takes 3 minutes to burn the DVD A and 4 minutes for B, but can work only for 24 hours and 10 minutes per series.

The packaging machine takes 5 minutes for the DVD A and 3 minutes for B, but it can work only for 25 hours in a row. Each DVD A is sold 50 Yuans and each DVD B is sold 40 Yuans.

- 1. Write the system of inequalities corresponding to this production.
- 2. Draw the lines corresponding to the production of each machine.
- 3. Shade the area corresponding to these conditions of production.
- 4. Write the equation corresponding to the total amount sold for this production.
- 5. Find the maximum number of DVD A and B which can be produced.
- 6. Draw the line of the sales corresponding to that maximum production.




北京景山学校 Name:

Mathematics - Elective Pre-Calc. - Senior 1 – TEST 1 Nov.2 (60 min.) - p.2/3

II– Parabolas and Hyperbolas : [40 pts] $f(x) = -\frac{1}{4}x^2 - x + 3$ $g(x) = \frac{2x+12}{x+4}$


- 1. Draw carefully the graphs of the two functions in the same system of coordinates. Show the axis of symmetry of the Parabola and the asymptotes of the Hyperbola
- 2. Calculate and show the coordinates of intersections with the 0x and the 0y axes.
- 3. Solve the equation f(x) = g(x) to find the coordinates of the intersection points of the Parabola and the Hyperbola.
- 4. Shade the area of points (x;y) corresponding to the system of inequalities : $y \le f(x) & y \ge g(x)$



Mathematics - Elective Pre-Calc. - Senior 1 – TEST 1 Nov.2 (60 min.) - p.3/3

III – Associated Functions and Transformations : graph the following functions in the system below

$$f1(x) = \frac{1}{4}x^2 - |x| - 3$$
; $f2(x) = -\sqrt{(x-3)^2} + 4$; $f3(x) = -\sqrt{9+9x} + 6$



November 15, 2010