


Numerical Sequences 2.1

Problem I : Let f be the function defined by $f(x) = \frac{2x+3}{x+4}$ for $x \geq 0$.

Study the Sequence defined by the formula $u_n = f(n) = \frac{2n+3}{n+4}$ for every $n \in \mathbb{N}$.

- Graph the function f on $[0; +\infty[$ and draw the first terms of the sequence (u_n) . Indicate from the graph whether or not the sequence is :
 - Monotonous (if yes how) :
 - Bounded (if yes, what are the boundaries ?)
 - Does-it seem to have a limit (if yes which one is it?)?
- Prove that (u_n) is increasing
- Prove that (u_n) is bounded by 0 and 2.
- Find for which value of n we have : $2 - \varepsilon < u_n < 2$ with $\varepsilon = 10^{-2}$
- Prove that for any $n \geq 1$ we have $|u_n - 2| \leq \frac{5}{n}$. Conclusion ?



- From the graph we can tell :
 - (u_n) is increasing;
 - $0.75 \leq u_n \leq 2$;
 - $\lim(u_n) = 2$
- From the properties of the function f , $f(x) = \frac{2x+3}{x+4} = 2 - \frac{5}{x+4}$ we can tell that (u_n) is increasing because f is increasing on $[0; +\infty[$ and $u_n = f(n)$.
- Because f is increasing on $[0; +\infty[$, $f(0) = .75$ and $\lim f(x) = 2$ then for any Integer n , $0.75 \leq u_n \leq 2$.

$$d) \quad 2 - \varepsilon \leq u_n \leq 2 + \varepsilon \Leftrightarrow 2 - \varepsilon \leq 2 - \frac{5}{n+4} \leq 2 + \varepsilon \Leftrightarrow -\varepsilon \leq -\frac{5}{n+4} \leq \varepsilon \Leftrightarrow -\varepsilon \leq \frac{5}{n+4} \leq \varepsilon$$

$$\Leftrightarrow (\varepsilon > 0) \frac{n+4}{5} \geq \frac{1}{\varepsilon} \Leftrightarrow n \geq \frac{5}{\varepsilon} - 4 \Leftrightarrow (\text{for } \varepsilon = \frac{1}{100}) n \geq 496$$

$$e) \quad |u_n - 2| = \left| \frac{-5}{n+4} \right| = \frac{5}{n+4} \leq \frac{5}{n} \quad \text{and} \quad \lim_{n \rightarrow +\infty} \left(\frac{5}{n} \right) = 0 \quad \text{then} \quad \lim_{n \rightarrow +\infty} |u_n - 2| = 0$$

which means that $\lim_{n \rightarrow +\infty} u_n = 2$

Numerical Sequences 2.2

Problem II : Let f be the function defined by $f(x) = \frac{2x+3}{x+4}$ for $x \geq 0$.


Study of the sequence (v_n) defined by $v_{n+1} = f(v_n) = \frac{2v_n+3}{v_n+4}$; $n \geq 1$ and $v_0 = 4$.

1. Graph the function f on $[0 ; +\infty [$ and draw the first terms of the sequence (v_n) . Find the coordinates of the intersection of (C_f) with the first bisector ($y=x$)
Indicate from the graph whether or not the sequence is :
 - i. Monotonous (if yes how) :
 - ii. Bounded (if yes, what are the boundaries ?)
 - iii. Does-it seem to have a limit (if yes which one is it?)?

2. Let $w_n = \frac{v_n - 1}{v_n + 3}$ for any $n > 0$.

Show that the new sequence (w_n) is a geometric sequence :

1. Find its first term and its reason.
2. Find the expression of w_n directly in function of n .
3. Deduct the limit of w_n .
4. Find the expression of v_n in function of w_n
5. Find the limit of v_n
6. Check the result on your graph.



- 1) From the graph of the sequence we can tell that the sequence (v_n) is decreasing, bounded by 1 and 4 and seems to be converging towards 1 (abscissa of the interception of f with the 1st bisector).
- 2) (w_n) is a geometric sequence because the ratio of two consecutive terms is constant :

$$w_{n+1} = \frac{v_{n+1} - 1}{v_{n+1} + 3} = \frac{\frac{2v_n+3}{v_n+4} - 1}{\frac{2v_n+3}{v_n+4} + 3} = \frac{v_n - 1}{5v_n + 15} = \frac{1}{5} \cdot \frac{v_n - 1}{v_n + 3} = \frac{1}{5} w_n ; w_0 = \frac{v_0 - 1}{v_0 + 3} = \frac{4 - 1}{4 + 3} = \frac{3}{7} \text{ and the reason } q = \frac{1}{5}$$

$$\therefore w_n = \frac{3}{7} \left(\frac{1}{5} \right)^n \Rightarrow \lim_{n \rightarrow +\infty} w_n = 0 \quad (\because |q| < 1)$$

$$w_n = \frac{v_n - 1}{v_n + 3} \Leftrightarrow v_n = \frac{1 + 3w_n}{1 - w_n} \text{ then } \lim_{n \rightarrow +\infty} v_n = \frac{1 + 3 \times 0}{1 - 0} = 1$$

The interception point of the graph of f and of the 1st bisector ($y=x$) yields the equation

$$\frac{2x+3}{x+4} = x \text{ and } x > 0 \Leftrightarrow x^2 + 2x - 3 = 0 \text{ and } x > 0 \Leftrightarrow x = 1$$